

0-0

Chapter 10

Advance of Mercury's Perihelion

3	10.1 Joyous Excitement	10-1
4	10.2 Newton's Simple Harmonic Oscillator	10-5
5	10.3 Newton's Orbit Analysis	10-5
6	10.4 Effective Potential: Einstein.	10-7
7	10.5 Einstein's Orbit Analysis	10-8
8	10.6 Predict Mercury's Perihelion Advance	10-10
9	10.7 Compare Prediction with Observation	10-12
10	10.8 Advance of the Perihelia of the Inner Planets	10-12
11	10.9 Check the Standard of Time	10-14
12	10.10 References	10-15

- 13 • *What does "advance of the perihelion" mean?*
- 14 • *You say Newton does not predict any advance of Mercury's perihelion in*
15 *the absence of other planets. Why not?*
- 16 • *The advance of Mercury's perihelion is tiny. So why should we care?*
- 17 • *Why pick out Mercury? Doesn't the perihelion of every planet change*
18 *with Earth-time?*
- 19 • *You are always shouting at me to say whose time measures various*
20 *motions. Why are you so sloppy about time in analyzing Mercury's orbit?*

CHAPTER

10

Advance of Mercury’s Perihelion

Edmund Bertschinger & Edwin F. Taylor *

This discovery was, I believe, by far the strongest emotional experience in Einstein’s scientific life, perhaps in all his life. Nature had spoken to him. He had to be right. “For a few days, I was beside myself with joyous excitement.” Later, he told Fokker that his discovery had given him palpitations of the heart. What he told de Haas is even more profoundly significant: when he saw that his calculations agreed with the unexplained astronomical observations, he had the feeling that something actually snapped in him.

—Abraham Pais

10.1 ■ JOYOUS EXCITEMENT

Tiny effect; large significance.

“Perihelion precession”?

What discovery sent Einstein into “joyous excitement” in November 1915? It was his calculation showing that his brand new (not quite completed) theory of general relativity gave the correct value for one detail of the orbit of the planet Mercury that had not been previously explained, an effect with the technical name **precession of Mercury’s perihelion**.

Newton: Sun-Mercury perihelion fixed.

Mercury (and every other planet) circulates around the Sun in a not-quite-circular orbit. In this orbit it oscillates in and out radially while it circles tangentially. A full Newtonian analysis predicts an elliptical orbit. Newton tells us that if we consider only the interaction between Mercury and the Sun, then the time for one 360-degree trip around the Sun is *exactly* the same as the time for one in-and-out radial oscillation. Therefore the orbital point closest to the Sun, the so-called **perihelion**, stays in the same place; the elliptical orbit does not shift around with each revolution—according to Newton. You will begin by verifying his nonrelativistic prediction for the simple Sun-Mercury system.

However, observation shows that Mercury’s orbit does indeed change. The perihelion moves forward in the direction of rotation of Mercury; it *advances*

*Draft of Second Edition of *Exploring Black Holes: Introduction to General Relativity* Copyright © 2017 Edmund Bertschinger, Edwin F. Taylor, & John Archibald Wheeler. All rights reserved. This draft may be duplicated for personal and class use.

10-2 Chapter 10

Advance of Mercury's Perihelion




FIGURE 1 Exaggerated view of the advance, during one century, of Mercury’s perihelion (and aphelion). The figure shows two elliptical orbits. One of these orbits is the one that Mercury traces over and over again in the year, say, 1900. The other is the elliptical orbit that Mercury traces over and over again in the year, say, 2000. The two are shifted with respect to one another, a rotation called *the advance (or precession) of Mercury’s perihelion*. The unaccounted-for precession in one Earth-century is about 43 arcseconds, less than the thickness of a line in this figure.

Observation:
perihelion advances.

52 with each orbit (Figure 1). The long (“major”) axis of the ellipse rotates. We
53 call this rotation of the axis the **advance (or precession) of the**
54 **perihelion.**

55 The **aphelion** is the point of the orbit farthest from the Sun; it advances
56 at the same angular rate as the perihelion (Figure 1).

Newton: Influence
of other planets,
predicts most of the
perihelion advance . . .

57 Observation shows that the perihelion of Mercury precesses at the rate of
58 574 arcseconds (0.159 degree) *per Earth-century*. (One degree equals 3600
59 arcseconds.) Newton’s mechanics accounts for 531 seconds of arc of this
60 advance by computing the perturbing influence of the other planets. But a
61 stubborn 43 arcseconds (0.0119 degree) per Earth-century, called a **residual**,
62 remains after all these effects are accounted for. This residual (though not its
63 modern value) was computed from observations by Urbain Le Verrier as early
64 as 1859 and more accurately later by Simon Newcomb (Box 1). Le Verrier
65 attributed the residual in Mercury’s orbit to the presence of an unknown inner
66 planet, tentatively named Vulcan. We know now that there is no planet
67 Vulcan. (Sorry, Mr. Spock!)

. . . but leaves
a *residual*.

Box 1. Simon Newcomb



FIGURE 2 Simon Newcomb
 Born 12 March 1835, Wallace, Nova Scotia.
 Died 11 July 1909, Washington, D.C.
 (Photo courtesy of Yerkes Observatory)

astronomers were those compiled by Simon Newcomb and his collaborator George W. Hill.

By the age of five Newcomb was spending several hours a day making calculations, and before the age of seven was extracting cube roots by hand. He had little formal education but avidly explored many technical fields in the libraries of Washington, D. C. He discovered the *American Ephemeris and Nautical Almanac*, of which he said, "Its preparation seemed to me to embody the highest intellectual power to which man had ever attained."

Newcomb became a "computer" (a person who computes) in the American Nautical Almanac office and by stages rose to become its head. He spent the greater part of the rest of his life calculating the motions of bodies in the solar system from the best existing data. Newcomb collaborated with Q. M. W. Downing to inaugurate a worldwide system of astronomical constants, which was adopted by many countries in 1896 and officially by all countries in 1950.

From 1901 until 1959 and even later, the tables of locations of the planets (so-called **ephemerides**) used by most

The advance of the perihelion of Mercury computed by Einstein in 1914 would have been compared to entries in the tables of Simon Newcomb and his collaborator.

Einstein correctly predicts residual precession.

Method: Compare in-and-out time with round-and-round time for Mercury.

68 Newton's mechanics says that there should be *no residual* advance of the
 69 perihelion of Mercury's orbit and so cannot account for the 43 seconds of arc
 70 per Earth-century which, though tiny, is nevertheless too large to be ignored
 71 or blamed on observational error. But Einstein's general relativity accounted
 72 for the extra 43 arcseconds on the button. Result: joyous excitement!

73 **Preview, Newton:** This chapter begins with Newton's approximations
 74 that lead to his no-precession conclusion (in the absence of other planets).
 75 Mercury moves in a near-circular orbit; Newton calculates the time for one
 76 orbit. The approximation also describes the small radial in-and-out motion of
 77 Mercury as if it were a harmonic oscillator moving back and forth about a
 78 potential energy minimum (Figure 3). Newton calculates the time for one
 79 in-and-out radial oscillation and compares it with the time for one orbit. The
 80 orbital and radial oscillation *T*-values are exactly equal (according to Newton),
 81 provided one considers only the Mercury-Sun interaction. He concludes that
 82 Mercury circulates around once in the same time that it oscillates radially
 83 inward and back out again. The result is an elliptical orbit that closes on itself.
 84 In the absence of other planets, Mercury repeats this exact elliptical path
 85 forever—according to Newton.

86 **Preview, Einstein:** In contrast, our general relativity approximation
 87 shows that these two times—the orbital round-and-round and the radial
 88 in-and-out *T*-values—are *not quite equal*. The radial oscillation takes place
 89 more slowly, so that by the time Mercury returns to its inner limit, the

10-4 Chapter 10

Advance of Mercury's Perihelion




FIGURE 3 Newton's effective potential, equation (5) (heavy curve), on which we superimpose the parabolic potential of the simple harmonic oscillator (thin curve) with the shape given by equation (3). Near the minimum of the effective potential, the two curves closely conform to one another.

90 circular motion has carried it farther around the Sun than it was at the
 91 preceding minimum r -coordinate. From this difference Einstein reckons the
 92 residual angular rate of advance of Mercury's perihelion around the Sun and
 93 shows that this predicted difference is close to the observed residual advance.
 94 Now for the details.

Comment 1. Relaxed about Newton's time and coordinate T

95 In this chapter we speak freely about Newton's time or Einstein's change in
 96 global T -value, without worrying about which we are talking about. We get away
 97 with this sloppiness for two reasons: (1) All observations are made from Earth's
 98 surface. Every statement about time should in principle be followed by the
 99 phrase, "as observed on Earth." (2) For this system, the effects of spacetime
 100 curvature on the rates of local clocks are so small that all time or T -measures
 101 give essentially the same rate of precession, as summarized in Section 10.11.
 102

10.2.3 ■ NEWTON'S SIMPLE HARMONIC OSCILLATOR

104 *Assume radial oscillation is sinusoidal.*

105 Why does the planet oscillate in and out radially? Look at the effective
 106 potential in Newton's analysis of motion, the heavy line in Figure 3. This
 107 heavy line has a minimum, the location at which the planet can ride around at
 108 constant r -value, tracing out a circular orbit. But with a slightly higher
 109 energy, it not only moves tangentially, it also oscillates radially in and out, as
 110 shown by the two-headed arrow in Figure 3.

111 How long does it take for one in-and-out oscillation? That depends on the
 112 shape of the effective potential curve near the minimum shown in Figure 3.
 113 But if the amplitude of the oscillation is small, then the effective part of the
 114 curve is very close to this minimum, and we can use a well-known
 115 mathematical theorem: If a continuous, smooth curve has a local minimum,
 116 then near that minimum a parabola approximates this curve. Figure 3 shows
 117 such a parabola (thin curve) superimposed on the (heavy) effective potential
 118 curve. From the diagram it is apparent that the parabola is a good
 119 approximation of the potential, at least near that local minimum.

In-and-out motion
 in parabolic potential . . .
 . . . predicts simple
 harmonic motion.

120 From introductory Newtonian mechanics, we know how a particle moves
 121 in a parabolic potential. The motion is called **simple harmonic oscillation**,
 122 described by the following expression:

$$x = A \sin \omega t \tag{1}$$

123 Here A is the amplitude of the oscillation and ω (Greek lower case omega) tells
 124 us how rapidly the oscillation occurs in radians per unit time. The potential
 125 energy per unit mass, V/m , of a particle oscillating in a parabolic potential
 126 follows the formula

$$\frac{V}{m} = \frac{1}{2} \omega^2 x^2 \tag{2}$$

127 To find the rate of oscillation ω of the harmonic oscillator, take the second
 128 derivative with respect to x of both sides of (2).

$$\frac{d^2 (V/m)}{dx^2} = \omega^2 \tag{3}$$

10.3 ■ NEWTON'S ORBIT ANALYSIS

130 *Round and round vs. in and out*

131 The in-and-out radial oscillation of Mercury does not take place around $r = 0$
 132 but around the r -value of the effective potential minimum. What is the
 133 r -coordinate of this minimum (call it r_0)? Start with Newton's equation (23)
 134 in Section 8.4:

Newton's
 equilibrium r_0

$$\frac{1}{2} \left(\frac{dr}{dt} \right)^2 = \frac{E}{m} - \left(-\frac{M}{r} + \frac{L^2}{2m^2 r^2} \right) = \frac{E}{m} - \frac{V_L(r)}{m} \quad (\text{Newton}) \tag{4}$$

10-6 Chapter 10

Advance of Mercury's Perihelion

135 This equation defines the effective potential,

$$\frac{V_L(r)}{m} \equiv -\frac{M}{r} + \frac{L^2}{2m^2r^2} \quad (\text{Newton}) \quad (5)$$

136 To locate the minimum of this effective potential, set its derivative equal to
137 zero:

$$\frac{d(V_L/m)}{dr} = \frac{M}{r^2} - \frac{L^2}{m^2r^3} = 0 \quad (\text{Newton}) \quad (6)$$

138 Solve the right-hand equation to find r_0 , the r -value of the minimum:

$$r_0 = \frac{L^2}{Mm^2} \quad (\text{Newton, equilibrium radius}) \quad (7)$$

Newton: In-and-out
time equals round-
and-round time.

139 We want to compare the rate ω_r of in-and-out radial motion of Mercury with
140 its rate ω_ϕ of round-and-round tangential motion. Use Newton's definition of
141 angular momentum, with increment dt of Newton's universal time, similar to
142 equation (10) of Section 8.2:

$$\frac{L}{m} \equiv r^2 \frac{d\phi}{dt} = r^2 \omega_\phi \quad (\text{Newton}) \quad (8)$$

143 where $\omega_\phi \equiv d\phi/dt$. Equation (8) gives us the angular velocity of Mercury along
144 its almost-circular orbit.

145 Queries 1 and 2 show that for Newton the radial in-and-out angular
146 velocity ω_r is equal to the orbital angular velocity ω_ϕ .

QUERY 1. Newton's angular velocity ω_ϕ of Mercury in orbit.

Set $r = r_0$ in (8) and substitute the result into (7). Show that at the equilibrium radius, $\omega_\phi^2 = M/r_0^3$ for Newton.

QUERY 2. Newton's radial oscillation rate ω_r for Mercury's orbit

We want to use (3) to find the angular rate of radial oscillation. Accordingly, take the second derivative of V_L in (5) with respect to r . Set $r = r_0$ in the resulting expression and substitute your value for L^2 in (7). Use (3) to show that at Mercury's orbital radius, $\omega_r^2 = M/r_0^3$, according to Newton.

158 **Important result:** *For Newton, Mercury's perihelion does not advance*
159 *when one considers only the gravitational interaction between Mercury and the*
160 *Sun.*

10.4. EFFECTIVE POTENTIAL: EINSTEIN

162 *Extra effective potential term advances perihelion.*

163 Now we repeat the analysis of radial and tangential orbital motion for the
 164 general relativistic case. Chapter 9 predicts the radial motion of an orbiting
 165 satellite. Multiply equations (4) and (5) of Section 9.1 through by 1/2 to
 166 obtain an equation similar to (4) above for the Newton's case:

$$\begin{aligned} \frac{1}{2} \left(\frac{dr}{d\tau} \right)^2 &= \frac{1}{2} \left(\frac{E}{m} \right)^2 - \frac{1}{2} \left(1 - \frac{2M}{r} \right) \left(1 + \frac{L^2}{m^2 r^2} \right) \quad (9) \\ &= \frac{1}{2} \left(\frac{E}{m} \right)^2 - \frac{1}{2} \left(\frac{V_L(r)}{m} \right)^2 \quad (\text{Einstein}) \end{aligned}$$

Set up general
relativity effective
potential.

167 Equations (4) and (9) are of similar form, and we use this similarity to make a
 168 general relativistic analysis of the harmonic radial motion of Mercury in orbit.
 169 In this process we adopt the *algebraic manipulations* of Newton's analysis in
 170 Sections 10.2 and 10.3 but apply them to the general relativistic expression (9).

Different time rates
of different clocks
do not matter.

171 Before we proceed, note three characteristics of equation (9). First, $d\tau$ on
 172 the left side of (9) is the differential wristwatch time $d\tau$, not the differential dt
 173 of Newton's universal time t . This different reference time is not necessarily
 174 fatal, since we have not yet decided which relativistic measure of time should
 175 replace Newton's universal time t . You will show in Section 10.11 that for
 176 Mercury the choice of which time to use (wristwatch time, global map
 177 T -coordinate, or even shell time at the r -value of the orbit) makes a negligible
 178 difference in our predictions about the rate of advance of the perihelion.

179 Note, second, that in equation (9) the relativistic expression $(E/m)^2$
 180 stands in the place of the Newtonian expression E/m in (4). However, both
 181 are constant quantities, which is all that matters in the analysis.

182 Evidence that we are on the right track results when we multiply out the
 183 second term of the first line of (9), which is the square of the effective
 184 potential, equation (18) of Section 8.4, with the factor one-half. Note that we
 185 have assigned the symbol $(1/2)(V_L/m)^2$ to this second term.

$$\begin{aligned} \frac{1}{2} \left(\frac{V_L(r)}{m} \right)^2 &= \frac{1}{2} \left(1 - \frac{2M}{r} \right) \left(1 + \frac{L^2}{m^2 r^2} \right) \quad (\text{Einstein}) \quad (10) \\ &= \frac{1}{2} - \frac{M}{r} + \frac{L^2}{2m^2 r^2} - \frac{ML^2}{m^2 r^3} \end{aligned}$$

Details of relativistic
effective potential

186 The heavy curve in Figure 4 plots this function. The second line in (10)
 187 contains the two effective potential terms that made up the Newtonian
 188 expression (5). The final term on the right of the second line of (10) describes
 189 an added attractive potential from general relativity. For the Sun-Mercury
 190 case at the r -value of Mercury's orbit, this term leads to the slight precession
 191 of the elliptical orbit. As r becomes small, the r^3 in the denominator causes
 192 this term to overwhelm all other terms in (10), which results in the downward
 193 plunge in the effective potential at the left side of Figure 4.

10-8 Chapter 10

Advance of Mercury's Perihelion




FIGURE 4 General-relativistic effective potential $(V_L/m)^2/2$ (heavy curve) and its approximation at the local minimum by a parabola (light curve) in order to analyse the radial excursion (double-headed arrow) of Mercury as simple harmonic motion. The effective potential curve is for a black hole, not for the Sun, whose effective potential near the potential minimum would be indistinguishable from the Newton's effective potential on the scale of this diagram. However, this minute difference accounts for the tiny residual precession of Mercury's orbit.

194 Finally, note third that the last term $(1/2)(V_L/m)^2$ in relativistic equation
 195 (9) takes the place of the Newton's effective potential V_L/m in equation (4).

196 In summary, we can manipulate general relativistic expressions (9) and
 197 (10) in nearly the same way that we manipulated Newton's expressions (4) and
 198 (5) in order to analyze the radial component of Mercury's motion and small
 199 perturbations of Mercury's elliptical orbit brought about by general relativity.

10.5 ■ EINSTEIN'S ORBIT ANALYSIS

201 *Einstein tweaks Newton's solution.*

202 Now analyze the radial oscillation of Mercury's orbit according to Einstein.

QUERY 3. Local minimum of Einstein's effective potential

Take the first derivative of the squared effective potential (10) with respect to r , that is find $d[(1/2)(V_L/m)^2]/dr$. Set this first derivative aside for use in Query 4. As a separate calculation, equate

this derivative to zero; set $r = r_0$, and solve the resulting equation for the unknown quantity $(L/m)^2$ in terms of the known quantities M and r_0 .

QUERY 4. Einstein's radial oscillation rate ω_r for Mercury in orbit.

We want to use (3) to find the rate of oscillation ω_r in the radial direction.

- A. Take the second derivative of $(1/2)(V_L/m)^2$ from (10) with respect to r . Set the resulting $r = r_0$ and substitute the expression for $(L/m)^2$ from Query 3 to obtain

$$\left[\frac{d^2}{dr^2} \left(\frac{1}{2} \frac{V_L^2}{m^2} \right) \right]_{r=r_0} = \omega_r^2 = \frac{M}{r_0^3} \frac{\left(1 - \frac{6M}{r_0} \right)}{\left(1 - \frac{3M}{r_0} \right)} \quad \text{(Einstein)} \quad (11)$$

$$\approx \frac{M}{r_0^3} \left(1 - \frac{6M}{r_0} \right) \left(1 + \frac{3M}{r_0} \right) \quad (12)$$

$$\approx \frac{M}{r_0^3} \left(1 - \frac{3M}{r_0} \right) \quad (13)$$

where we have made repeated use of the approximation inside the front cover in order to find a result to first order in the fraction M/r .

- B. For our Sun, $M \approx 1.5 \times 10^3$ meters, while for Mercury's orbit $r_0 \approx 6 \times 10^{10}$ meters. Does the value of M/r_0 justify the approximations in equations (12) and (13)?

Note that the coefficient M/r_0^3 in these three equations equals Newton's expression for ω_r^2 derived in Query 1.

Now compare ω_r , the in-and-out oscillation of Mercury's orbital r -coordinate with the angular rate ω_ϕ with which Mercury moves tangentially in its orbit. The rate of change of azimuth ϕ springs from the definition of angular momentum in equation (10) in Section 8.2:

$$\frac{L}{m} = r^2 \frac{d\phi}{d\tau} \quad \text{(Einstein)} \quad (14)$$

Note the differential wristwatch time $d\tau$ for the planet.

QUERY 5. Einstein's angular velocity

Square both sides of (14) and use your result from Query 3 to eliminate L^2 from the resulting equation. Show that at the equilibrium r_0 the result can be written

10-10 Chapter 10

Advance of Mercury's Perihelion

$$\omega_\phi^2 \equiv \left(\frac{d\phi}{d\tau}\right)^2 = \frac{M}{r_0^3} \left(1 - \frac{3M}{r_0}\right)^{-1} \quad (\text{Einstein}) \quad (15)$$

$$\approx \frac{M}{r_0^3} \left(1 + \frac{3M}{r_0}\right) \quad (16)$$

where again we use our approximation inside the front cover. Compare this result with equation (13) and with Newton's result in Query 1.

10.6. ■ PREDICT MERCURY'S PERIHELION ADVANCE

235 *Simple outcome, profound consequences*

Einstein: in-out rate differs from circulation rate.

236 According to Einstein, the advance of Mercury's perihelion springs from the
 237 difference between the frequency with which the planet sweeps around in its
 238 orbit and the frequency with which it oscillates in and out in r . In Newton's
 239 analysis these two frequencies are equal (for the interaction between Mercury
 240 and the Sun). But Einstein's theory shows that these two frequencies are
 241 *slightly* different; Mercury reaches its minimum r (its perihelion) at an
 242 incrementally greater angular position in each successive orbit. *Result:* the
 243 advance of Mercury's perihelion. In this section we compare Einstein's
 244 prediction with observation. But first we need to define what we are
 245 calculating.

246 What do we mean by the phrase "the period of a planet's orbit"? The
 247 period with respect to what? Here we choose what is technically called the
 248 **synodic period** of a planet, defined as follows:

Definition: synodic period

249 **DEFINITION 1. Synodic period of a planet**

250 The **synodic period** of a planet is the lapse in time (Newton) or lapse in
 251 global T -value (Einstein) for the planet to revolve once around the Sun
 252 with respect to the fixed stars.

253 **Comment 2. Fixed stars?**

"Fixed" stars?

254 What are the "fixed stars"? Chapter 14 The Expanding Universe shows that
 255 stars are anything but fixed. With respect to our Sun, stars move! However, stars
 256 that we now know to be very distant do not change angle rapidly from our point
 257 of view. Over a few hundred years—the lifetime of the field of astronomy
 258 itself—these stars may be called *fixed*.

259 The value T_r to make a complete in-and-out radial oscillation is

$$T_r \equiv \frac{2\pi}{\omega_r} \quad (\text{period of radial oscillation}) \quad (17)$$

260 In global coordinate lapse T_r , Mercury goes around the Sun, completing an
 261 angle

$$\omega_\phi T_r = \frac{2\pi\omega_\phi}{\omega_r} = (\text{Mercury revolution angle in } T_r) \tag{18}$$

262 which exceeds one complete revolution in radians by:

$$\omega_\phi T_r - 2\pi = T_r (\omega_\phi - \omega_r) = (\text{excess angle per revolution}) \tag{19}$$

QUERY 6. Difference in Einstein's oscillation rates

The two angular rates ω_ϕ and ω_r are *almost* identical in value, even in the Einstein analysis. Therefore we can write approximately:

$$\omega_\phi^2 - \omega_r^2 = (\omega_\phi + \omega_r)(\omega_\phi - \omega_r) \approx 2\omega_\phi(\omega_\phi - \omega_r) \tag{20}$$

A. Substitute equations (13) and (16) into the left side of (20):

$$\omega_\phi^2 - \omega_r^2 \approx \frac{M}{r_0^3} \left[\left(1 + \frac{3M}{r_0}\right) - \left(1 - \frac{3M}{r_0}\right) \right] = \frac{M}{r_0^3} \frac{6M}{r_0} \tag{21}$$

B. Equation (20) becomes:

$$\omega_\phi^2 - \omega_r^2 \approx \frac{M}{r_0^3} \frac{6M}{r_0} \approx \omega_\phi^2 \frac{6M}{r_0} \approx 2\omega_\phi(\omega_\phi - \omega_r) \tag{22}$$

C. Simplify the right-hand equation in (22), write the result as:

$$\omega_\phi - \omega_r \approx \frac{3M}{r_0} \omega_\phi \quad (\text{angular rates, Einstein}) \tag{23}$$

Equation (23) shows the difference in angular velocity between the tangential motion and the radial oscillation. From this rate difference we will calculate the advance of the perihelion of Mercury in one Earth-century.

Comment 3. What is X?

Symbols ω in (23) express rotation rates in radians per unit of—what? *Question:* What is X in the denominator of $d\phi/dX \equiv \omega$? Does X equal global coordinate T ? planet wristwatch time τ ? shell time t_{shell} at the average r -value of the orbit? *Answer:* It does not matter which of these quantities X represents, as long as this measure is the *same* on both sides of any resulting equation. Comment 1 told us to be relaxed about time. In the following Queries you use (23) to calculate the precession rate of Mercury in radians/second, then to convert this result to arcseconds/Earth-century.

10-12 Chapter 10

Advance of Mercury's Perihelion

10.7. ■ COMPARE PREDICTION WITH OBSERVATION

284 *Check out Einstein!*

285 Now compare our approximate relativistic prediction with observation.

QUERY 7. Mercury's angular velocity

The synodic period of Mercury's orbit is 7.602×10^6 seconds. To one significant digit, $\omega_\phi \approx 8 \times 10^{-7}$ radian/second. What is its value to three significant digits?

QUERY 8. Calculated coefficient

The mass M of the Sun is 1.477×10^3 meters and r_0 of Mercury's orbit is 5.80×10^{10} meters. To one significant digit, the coefficient $3M/r_0$ in (23) is 1×10^{-7} . Find this result to three significant digits.

QUERY 9. Advance of Mercury's perihelion in radians/second

From equation (23) and results of Queries 7 and 8, derive a numerical prediction of the advance of the perihelion of Mercury's orbit in radians/second. To one significant digit the result is 6×10^{-14} radians/second. Find the result to three significant digits.

QUERY 10. Advance of Mercury's perihelion in arcseconds per Earth-century.

Estimate the general relativity prediction of advance of Mercury's perihelion in arcseconds per century. Use results from preceding queries plus conversion factors inside the front cover plus the definition that 3600 arcseconds equals one degree. To one significant digit, the answer is 40 arcseconds/century. Find the result to three significant digits.

Observation and careful calculation agree.

310 A more accurate relativistic analysis predicts 42.980 arcseconds (0.011939
311 degrees) per Earth-century (Table 10.1). The observed rate of advance of the
312 perihelion is in perfect agreement with this value: 42.98 ± 0.1 arcseconds per
313 Earth-century. By what percentage did your prediction differ from
314 observation?

All planet orbits precess.

10.8. ■ ADVANCE OF THE PERIHELIA OF THE INNER PLANETS

316 *Help from a supercomputer.*

317 Do the *perihelia* (plural of *perihelion*) of other planets in the solar system also
318 advance as described by general relativity? Yes, but these planets are farther
319 from the Sun, and their orbits are less eccentric, so the magnitude of the
320 predicted advance is less than that for Mercury. In this section we compare our

Section 10.8 Advance of the Perihelia of the Inner Planets **10-13**

TABLE 10.1 Advance of the perihelia of the inner planets

Planet	Advance of perihelion in seconds of arc per Earth-century (JPL calculation)	r -value of orbit in AU*	Period of orbit in years
Mercury	42.980 ± 0.001	0.38710	0.24085
Venus	8.618 ± 0.041	0.72333	0.61521
Earth	3.846 ± 0.012	1.00000	1.00000
Mars	1.351 ± 0.001	1.52368	1.88089

*Astronomical Unit (AU): average r -value of Earth's orbit; inside front cover.

Computer analysis of precessions.

JPL multi-program computation.

321 estimated advance of the perihelia of the inner planets Mercury, Venus, Earth,
322 and Mars with results of an accurate calculation.

323 The Jet Propulsion Laboratory (JPL) in Pasadena, California, supports
324 an active effort to improve our knowledge of the positions and velocities of the
325 major bodies in the solar system. For the major planets and the moon, JPL
326 maintains a database and set of computer programs known as the Solar System
327 Data Processing System. The input database contains the observational data
328 measurements for current locations of the planets. Working together, more
329 than 100 interrelated computer programs use these data and the relativistic
330 laws of motion to compute locations of planets in the past and the future. The
331 equations of motion take into account not only the gravitational interaction
332 between each planet and the Sun but also interactions among all planets,
333 Earth's moon, and 300 of the most massive asteroids, as well as interactions
334 between Earth and Moon due to nonsphericity and tidal effects.

335 To help us with our project on perihelion advance, Myles Standish,
336 Principal Member of the Technical Staff at JPL, kindly used the numerical
337 integration program of the Solar System Data Processing System to calculate
338 orbits of the four inner planets over four centuries, from A.D. 1800 to A.D.
339 2200. In an overnight run he carried out this calculation twice, first with the
340 full program including relativistic effects and second "with relativity turned
341 off." Standish "turned off relativity" by setting the speed of light to 10^{10} times
342 its measured value, making light speed effectively infinite.

343 For each of the two runs, the perihelia of the four inner planets were
344 computed for the four centuries. The results from the nonrelativistic run were
345 subtracted from those of the relativistic run, revealing advances of the
346 perihelia per Earth-century accounted for only by general relativity. The
347 second column of Table 10.1 shows the results, together with the estimated
348 computational error.

QUERY 11. Approximate advances of the perihelia of the inner planets

Compare the JPL-computed advances of the perihelia of Venus, Earth, and Mars in Table 10.1 with approximate results calculated using equation (23).

10-14 Chapter 10

Advance of Mercury's Perihelion

10.9. CHECK THE STANDARD OF TIME

355 *Whose clock?*

356 We have been casual about whose time tracks the advance of the perihelion of
 357 Mercury and other planets; we even treated the global T -coordinate as a time,
 358 which is against our usual rules. Does this invalidate our approximations?

359

QUERY 12. Difference between shell time and Mercury's wristwatch time.

Use special relativity to find the fractional difference between planet Mercury's wristwatch time increment $\Delta\tau$ and the time increment Δt_{shell} read on shell clocks at the same average r_0 at which Mercury moves in its orbit at the average velocity 4.8×10^4 meters/second. By what fraction does a change of time from $\Delta\tau$ to Δt_{shell} change the total angle covered in the orbital motion of Mercury in one century? Therefore by what fraction does it change the predicted angle of advance of the perihelion in that century?

366

367

368

QUERY 13. Difference between shell time and global rain map T .

Find the fractional difference between shell time increment Δt_{shell} at r_0 and global map increment ΔT for r_0 equal to the average r -value of the orbit of Mercury. By what fraction does a change from Δt_{shell} to a lapse in global T alter the predicted angle of advance of the perihelion in that century?

372

374

QUERY 14. Does the time standard matter?

From your results in Queries 12 and 13, say whether or not the choice of a time standard—wristwatch time of Mercury, shell time, or map t —makes a detectable difference in the numerical prediction of the advance of the perihelion of Mercury in one Earth-century. Would your answer differ if the time were measured with clocks on Earth's surface?

380

381 DEEP INSIGHTS FROM MORE THAN THREE CENTURIES AGO

382 *Newton himself was better aware of the weaknesses inherent in his*
 383 *intellectual edifice than the generations that followed him. This fact*
 384 *has always roused my admiration.*

385 —Albert Einstein

386 We agree with Einstein. In the following quote from the end of his great work
 387 *Principia*, Isaac Newton summarizes what he knows about gravity and what
 388 he does not know. We find breathtaking the scope of what Newton says—and
 389 the integrity with which he refuses to say what he does not know. In the
 390 following, “feign” means “invent,” and since Newton's time “experimental
 391 philosophy” has come to mean “physics.”

392 “I do not ‘feign’ hypotheses.”

393 *Thus far I have explained the phenomena of the heavens and of our*
 394 *sea by the force of gravity, but I have not yet assigned a cause to*
 395 *gravity. Indeed, this force arises from some cause that penetrates as*
 396 *far as the centers of the sun and planets without any diminution of*
 397 *its power to act, and that acts not in proportion to the quantity of*
 398 *the surfaces of the particles on which it acts (as mechanical causes*
 399 *are wont to do) but in proportion to the quantity of solid matter,*
 400 *and whose action is extended everywhere to immense distances,*
 401 *always decreasing as the squares of the distances. Gravity toward*
 402 *the sun is compounded of the gravities toward the individual*
 403 *particles of the sun, and at increasing distances from the sun*
 404 *decreases exactly as the squares of the distances as far as the orbit*
 405 *of Saturn, as is manifest from the fact that the aphelia of the*
 406 *planets are at rest, and even as far as the farthest aphelia of the*
 407 *comets, provided that those aphelia are at rest. I have not as yet*
 408 *been able to deduce from phenomena the reason for these properties*
 409 *of gravity, and I do not “feign” hypotheses. For whatever is not*
 410 *deduced from the phenomena must be called a hypothesis; and*
 411 *hypotheses, whether metaphysical or physical, or based on occult*
 412 *qualities, or mechanical, have no place in experimental philosophy.*
 413 *In this experimental philosophy, propositions are deduced from the*
 414 *phenomena and are made general by induction. The*
 415 *impenetrability, mobility, and impetus of bodies, and the laws of*
 416 *motion and the law of gravity have been found by this method. And*
 417 *it is enough that gravity really exists and acts according to the laws*
 418 *that we have set forth and is sufficient to explain all the motions of*
 419 *the heavenly bodies and of our sea.*

420

—Isaac Newton

10.10 ■ REFERENCES

422 Initial quote: Abraham Pais, *Subtle Is the Lord: The Science and the Life of*
 423 *Albert Einstein*, Oxford University Press, New York, 1982, page 253.

424 Einstein’s analysis of precession of Mercury’s orbit in *Sitzungsberichte der*
 425 *Preussischen Akademie der Wissenschaften zu Berlin*, Volume 11, pages
 426 831–839 (1915). English translation by Brian Doyle in *A Source Book in*
 427 *Astronomy and Astrophysics, 1900–1975*, pages 820–825. Our analysis
 428 influenced by that of Robert M. Wald (*General Relativity*, University of
 429 Chicago Press, 1984, pages 142–143)

430 Myles Standish of the Jet Propulsion Laboratory ran the programs on the
 431 inner planets presented in Section 10. He also made useful comments on the
 432 project as a whole for the first edition.

10-16 Chapter 10

Advance of Mercury's Perihelion

433 Precession in binary neutron star systems: For B1913+16: J. M. Weisberg and
434 J. H. Taylor, *ApJ*, 576, 942 (2002). For J0737-3039: R. P. Breton et al,
435 *Science*, Vol. 321, 104 (2008)
436 Final Newton quote from I. Bernard Cohen and Anne Whitman, *Isaac*
437 *Newton, the Principia, A New Translation*, University of California Press,
438 Berkeley CA, 1999, page 943

439 Download File Name: Ch10AdvanceOfMercurysPerihelion160401v1.pdf