

Chapter 11 Orbits of Light

2	-
3	11.1 Turn a Stone into a Light Flash 11-1
4	11.2 Impact Parameter b 11-3
5	11.3 Equations of Motion for Light 11-5
6	11.4 Effective Potential for Light 11-9
7	11.5 Turning Points 11-12
8	11.6 Starlight Orbit: From Star to Observer 11-16
9	11.7 Integrate the Starlight Orbit 11-18
10	11.8 Multiple Starlight Orbits from every Star 11-19
11	11.9 Exercises 11-25
12	11.10 References 11-28

- 13 • *What variety of orbits does light follow around a black hole?*
- 14 • *Can a black hole reverse the direction of a light flash?*
- 15 • *Can light go into a circular orbit around a black hole? if so, is this*
- 16 *circular orbit stable?*
- 17 • *How many different orbits can light take from a single star to my eye?*

CHAPTER

11

Orbits of Light

Edmund Bertschinger & Edwin F. Taylor *

Then the sun god Ra emerged out of primal chaos.

—Egyptian creation story

And at once Kiho made his eyes to glow with flame—and the darkness became light.

—Tuamotuan (Polynesian) creation story

And God said, Let there be light: and there was light.

—first Biblical act of creation, Genesis 1:3

He bringeth them out of darkness unto light by His decree . . .

—Qur’an 5:16

Along with death came the Sun the Moon and the stars . . .

—Inuit creation story

11.1 ■ TURN A STONE INTO A LIGHT FLASH

Faster and faster, less and less mass

So far, observers are blind.

Thus far in this book almost all observers have been blind. Chapter 5 defined the shell observer but did not predict what he sees when he looks at stars or other objects outside his local inertial frame. The rain diver as she descends to the singularity (Chapter 7) peers in just two opposite directions—radially inward and radially outward. The explorer in her circular orbit around a black hole (Chapter 8) does not report what she sees—neither the starry heavens around her nor the black hole beneath her. In the present chapter we lay the groundwork to cure this blindness: we plot orbits of light in global map coordinates.

No local observation in this chapter

But this chapter still does not describe what any observer *sees*. Recall that we make every measurement and observation in a local inertial frame. The present chapter describes only map “starlight orbits,” for example the orbit

*Draft of Second Edition of *Exploring Black Holes: Introduction to General Relativity* Copyright © 2017 Edmund Bertschinger, Edwin F. Taylor, & John Archibald Wheeler. All rights reserved. This draft may be duplicated for personal and class use.

11-2 Chapter 11 Orbits of Light

46 that connects remote Star X with an observer at (or passing through) map
 47 location Y. The following Chapter 12 will tell us in what direction an observer
 48 at Y looks to see Star X.

Seeing is
 not believing.

49 What can we say about the global motion of light around, past, or into a
 50 spherically symmetric, nonspinning black hole? We ask here no small question:
 51 Almost every message from events in space comes to us by way of
 52 electromagnetic radiation of different frequencies. *Exceptions:* cosmic rays,
 53 neutrinos, and gravitational waves. A starlight orbit may deflect as it passes
 54 close to a massive object. Near a black hole this deflection can be radical;
 55 starlight can even go into a circular orbit. This and the following chapter make
 56 clear that for an observer near a black hole, seeing is definitely *not* believing!

Find orbits
 of light.

57 How do we plot the global orbit of light around a black hole? This is a
 58 new question; up until now we plotted light cones with short legs that sprout
 59 from a single event. Now we want to “connect the dots,” the events along an
 60 entire orbit of light that stretches from a specified distant star to a given local
 61 observer near a black hole.

Constant(s) of
 motion for light?

62 The *free stone* has two global constants of motion along its worldline: map
 63 energy E and map angular momentum L . Chapters 3 and 8 used the Principle
 64 of Maximal Aging to derive map expressions for each of these global constants
 65 of motion. Can we use the Principle of Maximal Aging to find constant(s) of
 66 motion for a light flash?

Principle of Maximal
 Aging does not apply
 directly to light.

67 The Principle of Maximal Aging says that a stone chooses a path across
 68 an adjoining tiny pair of segments along its worldline such that its wristwatch
 69 time is a maximum between a fixed initial event as the stone enters the pair
 70 and a fixed final event as it leaves the pair. But the Principle of Maximal
 71 Aging cannot apply directly to light, and for a fundamental reason: *The aging*
 72 *of a light flash along its worldline in a vacuum is automatically zero!* Aging
 73 $d\tau$ equals *zero* along every differential increment of the light flash worldline.
 74 *Question:* How can we possibly apply the Principle of Maximal Aging to light,
 75 whose aging is automatically zero?

Adapt Principle
 of Maximal Aging
 to light.

76 *Answer:* Sneak up on it! Start in flat spacetime far from a black hole.
 77 Think of a series of faster and faster stones, each stone with a smaller mass
 78 than the previous one. Let this series occur in such a way that the map energy
 79 E remains constant. Far from the black hole, map energy equals the
 80 measurable energy in a local inertial shell frame, in which the stone has
 81 squared speed v_{shell}^2 . Take the limit of equation (28) in Section 1.7 as $m \rightarrow 0$
 82 and $v_{\text{shell}} \rightarrow 1$:

$$E = \lim_{v_{\text{shell}} \rightarrow 1} \frac{m}{(1 - v_{\text{shell}}^2)^{1/2}} = \text{constant} \quad (\text{light, } r/M \gg 1) \quad (1)$$

Stone \rightarrow light
 as $m \rightarrow 0$
 and $v \rightarrow 1$

83 The present chapter analyzes consequences of this limit-taking process in (1).




FIGURE 1 Impact parameter b of a stone that approaches the black hole from a far away. Far from the black hole, we define b as the perpendicular offset between the line of motion of the approaching stone and the parallel line of motion of a test particle that makes a dive at constant ϕ into the black hole. Values of b and M determine whether or not the black hole captures the incoming stone.

11.2.4 IMPACT PARAMETER b

85 *Impact parameter from map angular momentum and map energy*

86 Chapter 8 analyzed circular orbits of a stone around the black hole. Now we
 87 want to describe more general orbits of both a stone and a light flash, so we
 88 define an orbit.

DEFINITION 1. Orbit: Stone or light flash

Definition:
orbit

90 An orbit is the worldline of a stone or light flash described by global
 91 coordinates. An orbit need not be circular around an origin, it need not
 92 be closed, it need not even remain in a bounded region of space.

93 A starlight orbit is a special case of the orbit:

DEFINITION 2. Starlight orbit

Definition:
starlight orbit

95 A starlight orbit is the orbit (Definition 1) of a light flash emitted by a star.

“Straight line”
verified in local
shell frame.

96 Think first about the orbit of a free stone far from the black hole—the
 97 right side of Figure 1. Far from the black hole this orbit is straight. How do we
 98 measure this orbit to verify that it is straight? As always, carry out
 99 measurements in a local inertial frame. We choose a shell frame (Section 5.7).
 100 Sufficiently far from the black hole this “local” shell frame can be quite large
 101 in the sense that over a significant range of r and ϕ special relativity correctly
 102 describes this orbit as a *straight line*. Now find a parallel straight line orbit
 103 that—by trial and error—moves without deflection to the center of the black
 104 hole (verified by measurement in a series of shell frames on both sides of
 105 Figure 1).

106 In a local inertial shell frame far from the black hole, we can measure
 107 perpendicular distances between parallel orbits. This leads to the definition of

11-4 Chapter 11 Orbits of Light

108 the **impact parameter**, with the symbol b . In a preliminary definition, we
 109 define the impact parameter of a stone far from the black hole:

**Preliminary
 definition:**
 impact parameter

110 **DEFINITION 3. Impact parameter b of a stone (preliminary)**
 111 The impact parameter b of a stone is the perpendicular
 112 distance—measured far from the black hole—between the straight orbit
 113 of the free stone and the parallel straight orbit of a second stone (test
 114 particle) that plunges at constant ϕ into the black hole.

QUERY 1. Every moving stone has an impact parameter

Show that every distant stone that changes global coordinates r or ϕ (or both) has an impact parameter—even a stone that moves away from the black hole.

120 Thus far the definition of the impact parameter is purely geometric.
 121 However, the right side of Figure 1 can be used to define angular momentum.
 122 The angular momentum of the stone takes the simple form:

$$L_{\text{far}} \equiv b_{\text{far}} p_{\text{far}} \quad (\text{stone in distant—flat—spacetime}) \quad (2)$$

Map angular
 momentum L

123 where p_{far} is the momentum of special relativity (Section 1.8). Equation (2)
 124 determines the value of L where $r/M \gg 1$, that is where spacetime is flat.
 125 However L is a map constant of motion, the same everywhere around the
 126 black hole. Therefore its value, calculated from (2) far from the black hole, is
 127 the same close to the black hole.

128 Recall equation (39) for a stone in Section 1.9, with p defined in (2):

$$m^2 = E^2 - p^2 = E^2 - \left(\frac{L}{b}\right)^2 \quad (\text{stone, flat spacetime}) \quad (3)$$

Impact parameter
 of a stone

129 Solve this equation for b , in which b and L are either both positive or both
 130 negative:

$$b \equiv \frac{L}{(E^2 - m^2)^{1/2}} \quad (\text{impact parameter for a stone, everywhere}) \quad (4)$$

131
 132 Both map energy E and map angular momentum L are map constants of
 133 motion and m is an invariant quantity. Therefore equation (4) is valid close to
 134 the black hole as well as far away. Even though it was derived assuming flat
 135 spacetime, we take (4) to define b everywhere. Close to the black hole, b is no
 136 longer the perpendicular distance of Definition 3. But every orbit has an L and
 137 an E and therefore can be assigned a unique value of b .

138 For light, carry out the limit-taking process demanded in (1), with
 139 constant E but decreasing m . The limit $m \rightarrow 0$ defines the impact parameter
 140 for light:

Impact parameter
 of a light flash

Section 11.3 Equations of Motion for Light **11-5**

$$b \equiv \frac{L}{E} \quad (\text{impact parameter of light, everywhere}) \quad (5)$$

141

142 This leads to the final definition of the impact parameter for a stone or a
143 light flash around a black hole:

144

DEFINITION 4. Impact parameter b

Definition:
impact parameter b

145

The **impact parameter** b for a stone is given by (4) and for a light flash
146 by (5).

146



147

148

149

150

Objection 1. *You use two perfectly good constants of motion, L and E and give a geometric interpretation for a combination of them. So what? I can define a thousand combinations of L and E . Who cares? I didn't need any such combination for a stone. Why are you wasting my time?*



151

152

153

154

155

We introduce b because neither L alone or E alone will be helpful when $m \rightarrow 0$. Equations of motion for light derived below depend only on the fraction L/E and no other combination. Global motion of a stone depends on two constants of motion, L and E . Global motion of light is simpler, completely described by one constant of motion, $b \equiv L/E$. Rejoice!

156

157

158

159

We have defined impact parameter, but we have not yet predicted the global motion of a light flash near the black hole. To obtain equations of motion for light, we again apply the limit-taking process of equation (1), in this case to the equations of motion for a stone from Chapter 8.

11.3 EQUATIONS OF MOTION FOR LIGHT

161

A single constant of motion for light, namely b

Flat starlight
wavefront approaching
the black hole . . .

162

163

164

165

Light spreads out from a star as a spherical wave. We assume that every star is so far away that as its starlight approaches our black hole—but still travels in flat spacetime—it forms a flat wavefront (right side of Figure 2).

. . . is equivalent to
a bundle of parallel
straight orbits.

166

167

168

169

170

We already have another powerful way to describe starlight in flat spacetime: as a bundle of parallel straight orbits. Figure 2 displays four starlight orbits from a single star, each with a different impact parameter b , as these orbits approach the black hole. Far from the black hole (right side of the figure) these starlight orbits remain parallel to one another. Close to the black hole (left side of the figure) they diverge: Only the orbit with $b/M = 0$ remains straight. Starlight Orbit 1 deflects but escapes; Starlight Orbit 2 enters a circular orbit; Starlight Orbit 3 plunges to the center of the black hole.

Close to the black
hole, orbits from
the star are neither
parallel nor straight.

171

172

173

174

Starlight Orbit 2 in Figure 2 is unique; it enters a circular orbit at $r = 3M$. We call this orbit *critical* and its impact parameter the *critical impact parameter*, b_{critical} . In Query 3 you show that the critical impact parameter has the value $b_{\text{critical}} = (27)^{1/2}M$.

Critical impact
parameter

175

176

11-6 Chapter 11 Orbits of Light




FIGURE 2 Jagged lines separate flat spacetime far from the black hole (on the right) from curved spacetime near the black hole (on the left). The right side of this plot shows two ways to visualize starlight orbits far from the black hole: first as a set of straight parallel orbits, second as a flat wavefront. On the left side of this plot, near the black hole, only the starlight orbit with $b/M = 0$ remains straight, while starlight orbits 1 through 3, originally parallel, diverge: Starlight Orbit 1 with the impact parameter $b/M = 7$ deflects but escapes. Starlight Orbit 2 with the so-called *critical impact parameter* b_{critical}/M , equation (28), becomes an unstable circular orbit at $r/M = 3$. Starlight Orbit 3 with $b/M = 4$ crosses the event horizon and ends at the singularity.

177 We need general equations of motion of light, which we now derive using
 178 the limiting process of equation (1). Start with equations of motion of a stone
 179 from Section 8.3, written in slightly altered form:

$$\frac{dr}{d\tau} = \pm \left[\left(\frac{E}{m} \right)^2 - \left(1 - \frac{2M}{r} \right) \left(1 + \frac{L^2}{m^2 r^2} \right) \right]^{1/2} \quad (\text{stone}) \quad (6)$$

$$\frac{d\phi}{d\tau} = \frac{L}{mr^2} \quad (\text{stone}) \quad (7)$$

$$\frac{d\tau}{dT} = \frac{\left(1 - \frac{2M}{r} \right)}{\frac{E}{m} \pm \left(\frac{2M}{r} \right)^{1/2} \left[\left(\frac{E}{m} \right)^2 - \left(1 - \frac{2M}{r} \right) \left(1 + \frac{L^2}{m^2 r^2} \right) \right]^{1/2}} \quad (8)$$

Comment 1. Choice of signs for the motion of a stone

180 We choose the stone's wristwatch time to advance as the stone moves along its
 181 worldline. Therefore the upper (+) sign in (6) is for a stone with increasing r and
 182 the lower (-) sign is for a stone with decreasing r . The \pm sign in the
 183 denominator of equation (8) has the same meaning.
 184

185 In order to describe the motion of light, we need to eliminate $d\tau$ from
 186 these equations, because adjacent events along the worldline of a light flash

Section 11.3 Equations of Motion for Light **11-7**

187 have zero wristwatch time lapse between them: $d\tau = 0$. Multiply both sides of
 188 (6) by the corresponding sides of (8), then factor out and cancel (E/m) from
 189 the resulting numerator and denominator.

$$\begin{aligned} \frac{dr}{dT} &= \frac{dr}{d\tau} \frac{d\tau}{dT} && \text{(stone)} && (9) \\ &= \pm \frac{\left(1 - \frac{2M}{r}\right) \left[1 - \left(\frac{m}{E}\right)^2 \left(1 - \frac{2M}{r}\right) \left(1 + \frac{L^2}{m^2 r^2}\right)\right]^{1/2}}{1 \pm \left(\frac{2M}{r}\right)^{1/2} \left[1 - \left(\frac{m}{E}\right)^2 \left(1 - \frac{2M}{r}\right) \left(1 + \frac{L^2}{m^2 r^2}\right)\right]^{1/2}} \end{aligned}$$

190 Equation (1) requires that for light $m \rightarrow 0$ while E remains constant. Apply
 191 these requirements to (9). The result is our first equation of motion for light:

$$\frac{dr}{dT} = \pm \frac{\left(1 - \frac{2M}{r}\right) \left[1 - \left(1 - \frac{2M}{r}\right) \left(\frac{L}{rE}\right)^2\right]^{1/2}}{1 \pm \left(\frac{2M}{r}\right)^{1/2} \left[1 - \left(1 - \frac{2M}{r}\right) \left(\frac{L}{rE}\right)^2\right]^{1/2}} \quad \text{(light) (10)}$$

192 Carry out a similar procedure on equations (7) and (8): multiply their
 193 corresponding sides $d\phi/dT = (d\phi/d\tau)(d\tau/dT)$, factor out E/m in the
 194 denominator, cancel m with one in the numerator, then let $m \rightarrow 0$. The result
 195 is our second equation of motion for light:

$$\frac{d\phi}{dT} = \frac{\frac{L}{r^2 E} \left(1 - \frac{2M}{r}\right)}{1 \pm \left(\frac{2M}{r}\right)^{1/2} \left[1 - \left(1 - \frac{2M}{r}\right) \left(\frac{L}{rE}\right)^2\right]^{1/2}} \quad \text{(light) (11)}$$

196 To construct our third equation of motion for light, combine (10) with (11):

$$\frac{dr}{d\phi} = \left(\frac{dr}{dT}\right) \left(\frac{dT}{d\phi}\right) = \pm \frac{r^2 E}{L} \left[1 - \left(1 - \frac{2M}{r}\right) \left(\frac{L}{rE}\right)^2\right]^{1/2} \quad \text{(light) (12)}$$

197 Equations (10) through (12) are the equations of motion for light. The choice
 198 of signs in these equations is the same as for a stone, given in Comment 1.

199 Our three equations of motion for light contain a wonderful surprise: The
 200 only quantity we need to describe the orbit of light is the ratio L/E . *Meaning:*
 201 The orbit of light near a black hole is completely determined by the single
 202 value of the ratio L/E instead of by the separate values of the map constants
 203 of motion L and E . And equation (5) tells us that this ratio equals the impact
 204 parameter for light.

205 Substitute the expression $b = L/E$ into equations (10) through (12):

Light motion depends
 on only $L/E = b$.

11-8 Chapter 11 Orbits of Light

$$\frac{dr}{dT} = \pm \frac{\left(1 - \frac{2M}{r}\right) \left[1 - \left(1 - \frac{2M}{r}\right) \left(\frac{b}{r}\right)^2\right]^{1/2}}{1 \pm \left(\frac{2M}{r}\right)^{1/2} \left[1 - \left(1 - \frac{2M}{r}\right) \left(\frac{b}{r}\right)^2\right]^{1/2}} \quad (\text{light}) \quad (13)$$

$$\frac{d\phi}{dT} = \frac{\frac{b}{r^2} \left(1 - \frac{2M}{r}\right)}{1 \pm \left(\frac{2M}{r}\right)^{1/2} \left[1 - \left(1 - \frac{2M}{r}\right) \left(\frac{b}{r}\right)^2\right]^{1/2}} \quad (\text{light}) \quad (14)$$

$$\frac{dr}{d\phi} = \pm \frac{r^2}{b} \left[1 - \left(1 - \frac{2M}{r}\right) \left(\frac{b}{r}\right)^2\right]^{1/2} \quad (\text{light}) \quad (15)$$

206 An identical square-bracket expression appears multiple times in these
 207 equations. To simplify them, define a new function $F(b, r)$:

$$F(b, r) \equiv \left[1 - \frac{b^2}{r^2} \left(1 - \frac{2M}{r}\right)\right]^{1/2} \quad (\text{light}) \quad (16)$$

Equations of
 motion for light

208
 209 so that equations of motion for light become:

$$\frac{dr}{dT} = \pm \frac{\left(1 - \frac{2M}{r}\right) F(b, r)}{1 \pm \left(\frac{2M}{r}\right)^{1/2} F(b, r)} \quad (\text{light}) \quad (17)$$

$$\frac{d\phi}{dT} = \frac{\frac{b}{r^2} \left(1 - \frac{2M}{r}\right)}{1 \pm \left(\frac{2M}{r}\right)^{1/2} F(b, r)} \quad (\text{light}) \quad (18)$$

$$\frac{dr}{d\phi} = \pm \frac{r^2}{b} F(b, r) \quad (\text{light}) \quad (19)$$

210
 211 The \pm signs in equations (17) through (19) have the same interpretation as in
 212 (6) through (8) and also (10) through (12), namely the upper (+) sign
 213 describes light with increasing r and the lower (-) describes light with
 214 decreasing r .

215 Chapters 9 and 10 use interactive software GRorbits to plot orbits of a
 216 stone. GRorbits also integrates equations (17) through (19) for light. Given

Section 11.4 Effective Potential for Light **11-9**

217 the value of b and initial location, the software plots the orbit and outputs a
 218 spreadsheet with global coordinates (T, r, ϕ) of events along the orbit.

219 Equations of motion for light look complicated. We now derive a simple
 220 way to visualize the global r -motion of light using the effective potential,
 221 modeled after the effective potential for a stone in Section 8.4.

11.4 ■ EFFECTIVE POTENTIAL FOR LIGHT

223 *Describe global motion of light at a glance.*

224 The present section sets up an effective potential for a light orbit in order to
 225 visualize its r -component of motion simply and directly. Recall equation (21)
 226 in Section 8.4 that relates the r -motion of a stone to its effective potential:

$$\left(\frac{dr}{d\tau}\right)^2 = \left(\frac{E}{m}\right)^2 - \left(\frac{V_L(r)}{m}\right)^2 \quad (\text{stone}) \quad (20)$$

227 The key idea of this equation is that the first term on the right is a constant of
 228 the stone's motion—independent of location—while the second term is a
 229 function of r —independent of the properties or motion of the stone. We
 230 defined the second term to be the effective potential for a stone.

231 To make similar predictions about the r -motion of light, we seek an
 232 equation with the same form as (20). To find this equation, square both sides
 233 of (17), rearrange the results, and multiply through by $(M/b)^2$ to obtain:

$$\left(\frac{M}{b}\right)^2 \left(1 - \frac{2M}{r}\right)^{-2} \left[1 \pm \left(\frac{2M}{r}\right)^{1/2} F(b, r)\right]^2 \left(\frac{dr}{dT}\right)^2 = \left(\frac{M}{b}\right)^2 F^2(b, r) \quad (21)$$

234 On the left side of (21) we define the function

$$A^2(b, r) \equiv \left(\frac{M}{b}\right)^2 \left(1 - \frac{2M}{r}\right)^{-2} \left[1 \pm \left(\frac{2M}{r}\right)^{1/2} F(b, r)\right]^2 \quad (\text{light}) \quad (22)$$

235 and on the right side of (21) we substitute for $F^2(b, r)$ from (16).

$$\left(\frac{M}{b}\right)^2 F^2(b, r) = \frac{M^2}{b^2} - \frac{M^2 b^2}{b^2 r^2} \left(1 - \frac{2M}{r}\right) \quad (\text{light}) \quad (23)$$

effective potential
for light

236 Substitute the left sides of (22) and (23) into (21) and write the result as:

11-10 Chapter 11 Orbits of Light

Box 1. Use of the effective potential for a stone and for a light flash

Compare and contrast the forms and uses of effective potentials for a stone and for a light flash:

$$\left(\frac{dr}{d\tau}\right)^2 = \left(\frac{E}{m}\right)^2 - \left(\frac{V_L(r)}{m}\right)^2 \quad (\text{stone}) \quad (26)$$

$$A^2 \left(\frac{dr}{dT}\right)^2 = \left(\frac{M}{b}\right)^2 - \left(\frac{V(r)}{M}\right)^2 \quad (\text{light}) \quad (27)$$

For a stone:

- V_L depends on both L and r .
- The turning point occurs where $V_L = \pm E$.
- $|E| < |V_L|$ is forbidden
- When $|E| \geq |V_L|$, equation (26) gives $|dr/d\tau|$ in terms of r, L, E .

For a light flash:

- V depends on r alone.
- The turning point occurs where $V/M = \pm M/b = \pm ME/L$, not E alone.
- $|M/b| < |V/M|$ is forbidden
- When $|M/b| \geq |V/M|$, equation (27) gives $|dr/dT|$ in terms of r, b .

What's the difference between the two cases?

For light, L has been removed from the effective potential and combined with E ; only $b = L/E$ remains. Impact parameter b can be taken completely out of the effective potential, so V depends only on r . This makes orbits of light *simpler* than orbits of a stone. Only *one* constant of motion is needed, not two.

$$A^2(b, r) \left(\frac{dr}{dT}\right)^2 = \left(\frac{M}{b}\right)^2 - \left(\frac{V(r)}{M}\right)^2 \quad (\text{light}) \quad (24)$$

where (25) defines the square of the **effective potential for light**

$$\left(\frac{V(r)}{M}\right)^2 \equiv \frac{M^2}{r^2} \left(1 - \frac{2M}{r}\right) \quad (\text{light}) \quad (25)$$

237

238 Figure 3 plots positive values of the effective potential for light. In Query 2
 239 you show that the coefficient $A^2(b, r)$ in equation (22) is well behaved when
 240 light descends to the event horizon, provided $b \neq 0$.

241 Box 1 compares and contrasts effective potentials for light and for stones.

242

QUERY 2. Approaching the event horizon

What happens to the left side of (24) as $r/M \rightarrow 2^+$, that is as light approaches the event horizon from above? Just above the event horizon set $r/M = 2(1 + \epsilon)$ where $0 < \epsilon \ll 1$ and use our standard approximation (inside the front cover) to show that coefficient $A^2(b, r)$ in (24) is well behaved even as light descends to the event horizon, provided $b \neq 0$.

248

Quick predictions with the effective potential

249 With the effective potential we can predict—at a glance—the r -component
 250 of light motion. The first term, $(M/b)^2$, on the right side of (24) is a constant
 251 of motion, the same everywhere along the orbit. The second term is a function
 252 of r and does not include b . Figure 3 and its caption also contain a preview of
 253 *turning points*, which we analyze more fully in Section 11.4.

Section 11.4 Effective Potential for Light **11-11**

254 *Huge payoff:* The right side of (24) does not include the energy or angular
 255 momentum of light. *One effective potential applies to light orbits of every*
 256 *energy and every angular momentum.* In particular, it applies to
 257 electromagnetic radiation of all wavelengths: radio waves; microwaves;
 258 infrared, visible, and ultraviolet light; X-rays; and gamma rays! (This result
 259 assumes that the wavelength of light is small compared with the coordinate
 260 separations over which spacetime curvature changes appreciably.)

Same effective potential for light of EVERY energy (EVERY wavelength)

QUERY 3. Critical impact parameter

- A. Show that the peak of the effective potential occurs at $r/M = 3$.
- B. Verify that the so-called **critical value** of the impact parameter at $r/M = 3$ is

$$\frac{b_{\text{critical}}}{M} = (27)^{1/2} = 5.196\ 152\ 42 \quad (\text{light, critical impact parameter}) \quad (28)$$

- C. From Figure 3 read off approximate values of b/M and r/M for the circular orbit. Compare these values with the analytic results of Items A and B.

Effective potentials reveals turning points.

Both the effective potential for light and effective potentials for stones enable us to find the r -coordinate at which the r -component of motion goes to zero, which occurs for a circular orbit and also at what we call a *turning point* (Section 8.4 and Section 11.5).

DEFINITION 5. Plunge Orbit, Bounce Orbit, Trapped Orbit

Figure 3 sorts all light orbits near a black hole into three categories, which we give names to simplify our analysis:

- **Plunge Orbit:** A plunge orbit is an incoming or outgoing orbit with $|b| < b_{\text{critical}}$ that passes above the peak of the effective potential curve in Figure 3. A starlight Plunge Orbit is—by definition—an incoming orbit that *plunges* through the event horizon to the singularity. Outside the event horizon light can, in principle, move in either direction along the plunge orbit shown. We call this a plunge orbit, whether r decreases or increases.
- **Bounce Orbit:** A bounce orbit is an incoming or outgoing orbit with $|b| > b_{\text{critical}}$. The bounce orbit exists only to the right of the effective potential in Figure 3 and below its peak. A starlight Bounce Orbit is—by definition—an orbit that initially moves inward, then reverses its r -component of motion—its r -coordinate *bounces*—at a turning point on the outer edge of the effective potential, while its ϕ -component of motion continues. After the bounce, the light moves outward on the same horizontal line in the figure, and escapes to infinity. A Bounce Orbit cannot reach the singularity.

Definitions:
 Plunge Orbit
 Bounce Orbit
 Trapped Orbit

11-12 Chapter 11 Orbits of Light




FIGURE 3 Examples of the three categories into which we sort all orbits (Definition 5). *Horizontal Line (1)*: a Plunge Orbit with $M/b = 1/4$ that enters the black hole. *Horizontal Line (2)*: the orbit with $M/b_{\text{critical}} = 1/(27)^{1/2}$ that reaches the peak of the effective potential—marked with a little filled circle—and enters an unstable circular orbit there. *Horizontal Line (3)*: a Bounce Orbit with $M/b = 1/7$ approaches the black hole, reverses its r -motion at the *outer turning point* (Section 11.6), and moves away from the black hole. The Trapped Orbit with $M/b = 1/7$ originates in the narrow horizontal region between the event horizon and the effective potential curve and moves inward through the event horizon.

293
294
295
296
297
298

- **Trapped Orbit:** A trapped orbit is an orbit with $|b| > b_{\text{critical}}$ to the left of the effective potential in Figure 3 and below its peak. No starlight orbit can be a Trapped Orbit. An initially outgoing Trapped Orbit outside the event horizon reverses its r -component of motion at the inner turning point on the inner edge of the effective potential. *Every* Trapped Orbit reaches the singularity unless intercepted.

299
300
301

The horizontal line for M/b_{critical} in Figure 3 is the dividing line between these different categories of orbits. Figure 4 shows Plunge and Bounce Orbits; Figure 5 shows two Trapped Orbits.

11.5.2 ■ TURNING POINTS

303
304
305
306

The r -motion of light can reverse at a turning point.

At a turning point the r -component of motion goes to zero, while the ϕ -component of motion continues. Little filled squares in Figures 3 through 5 mark what we call **outer and inner turning points**.

Section 11.5 Turning Points 11-13




FIGURE 4 Top two panels: Plunge Orbits. Bottom two panels: Bounce Orbits, each with a little filled square at the turning point (Section 11.4). Middle two panels: b -values straddle $b_{\text{critical}}/M = 5.19615\dots$, for which the orbit enters a knife-edge circular orbit.

DEFINITION 6. Turning Point

A turning point is the r -value at which the right side of equation (24) equals zero, where M/b equals the value of the effective potential.

- An **outer turning point** is to the right and below the peak of the effective potential (see Figure 3).
- An **inner turning point** is to the left and below this peak. The peak itself is the location of the unstable (knife-edge) circular orbit of light.
- A **circular orbit point** is the r -value at which the effective potential is maximum. This is the r -location of an unstable (knife-edge) circular orbit for light.

We use the subscript **tp** to label the r -coordinate of a turning point.

Example: In Figure 3, Orbit 3 with $|b/M| = 7$ reverses its r -motion at

Definitions:
 Turning point
 Outer turning point
 Inner turning point
 Circular orbit poin

Turning point
 subscript: tp

11-14 Chapter 11 Orbits of Light




FIGURE 5 Two Trapped Orbits that originate from the same point just outside the event horizon at $r/M = 2^+$ (little open circle). One orbit has $b/M = +6$ with an inner turning point (little filled square); the other has $b/M = -6$ and no turning point. Both orbits reach the singularity at $r/M = 0$. Figure 6 adds labels to this plot.

320 $r_{\text{tp}} = 5.617M$. Any outgoing light with $|b/M| = 7$ that arrives at the inner
 321 turning point at $r_{\text{tp, inner}} = 2.225M$ thereafter moves with $dr < 0$ and enters
 322 the black hole.

323 Equations (24) and (25) tell us that the turning point r_{tp} , the
 324 r -coordinate at which $dr/dT = 0$ and motion is purely tangential, occurs for
 325 the value of b given by:

$$b/M = \pm \frac{r_{\text{tp}}/M}{\left(1 - \frac{2M}{r_{\text{tp}}}\right)^{1/2}} \quad (\text{given } r_{\text{tp}}, \text{ find } b) \quad (29)$$

Comment 2. No turning point inside the event horizon

Turning points
 only for $b^2 > b_{\text{critical}}^2$

326 Equation (29) guarantees that there can be no turning point for light inside the
 327 event horizon, because b/M on the left side is necessarily a real quantity, while
 328 the right side of (29) is imaginary for $r_{\text{tp}} < 2M$.
 329

Derive r_{tp}
 from b .

330 Equation (29) gives us the value of b when we know the r -coordinate r_{tp} of the
 331 turning point. More often, we know the value of b and want to find the
 332 r -coordinate of the turning point. In that case, convert (29) into a cubic
 333 equation in r_{tp} :

$$r_{\text{tp}}^3 - b^2 r_{\text{tp}} + 2Mb^2 = 0 \quad (\text{given } b, \text{ find } r_{\text{tp}}) \quad (30)$$

QUERY 4. Optional: Some consequences of turning points.

A. From equations (24) and (25) show that a light orbit with a given value of b cannot exist in a range of r -coordinates determined by the following inequality:

$$r^3 - b^2 r + 2Mb^2 < 0 \quad (\text{region with no light orbits}) \quad (31)$$

Section 11.5 Turning Points **11-15**

B. Show that inequality (31) describes the shaded region under the effective potential curve in Figure 3. In other words, light cannot penetrate the effective potential curve.

Find the turning points

Equation (30) is cubic—includes a third power of r_{tp} . Cubic equations can be difficult to solve. Here are analytic solutions of (30). The first two yield r values of the outer and inner turning points, respectively, such as those in Figure 3. In Query 4 you show that the third solution is real but negative, so cannot represent the always-positive map r -coordinate:

$$r_{\text{tp}} = 3M \left[\frac{1}{2} - \cos(\psi - 120^\circ) \right]^{-1} \tag{32}$$

(Outer turning points lie at $r > 3M$.)

$$r_{\text{tp, inner}} = 3M \left[\frac{1}{2} - \cos(\psi + 120^\circ) \right]^{-1} \tag{33}$$

(Inner turning points lie between $r/M = 2$ and $r/M = 3$.)

$$r_{\text{NO}} = 3M \left[\frac{1}{2} - \cos \psi \right]^{-1} \tag{34}$$

(Yields negative r : not physical.)

For all three solutions, ψ depends on b as follows:

$$\psi \equiv \frac{1}{3} \arccos \left(\frac{54M^2}{b^2} - 1 \right) \quad (|b| \geq b_{\text{critical}}, 0 \leq \psi \leq \pi) \tag{35}$$

We take what is called the *principle value* of the arccos z , that is the angle between 0 and π radians whose cosine is z . Recall that the magnitude of the cosine is never greater than one. Therefore turning points exist only when the arccos function (35) exists, that is when $b^2 \geq b_{\text{critical}}^2$ or when the horizontal line for $(M/b)^2$ in Figure 3 is at or below the peak of the effective potential. This makes graphical, as well as analytic, sense.

QUERY 5. Unphysical third solution

Show that the third solution (34) yields a negative value for r , which cannot represent the non-negative r -coordinate.

QUERY 6. Examples of turning points

- A. For the outer and inner turning points of the orbit with $|b/M| = 7$, derive the numerical values $r_{\text{tp}} = 5.617M$ and $r_{\text{tp, inner}} = 2.225M$. Use Figure 3 to verify these r -coordinates approximately.
- B. Show that $F(\dot{t}_2 r) = 0$ at the turning points.

11-16 Chapter 11 Orbits of Light




FIGURE 6 Elaboration of Figure 5. Two Trapped Orbits originate from just outside the event horizon at $r_{\text{src}}/M = 2^+$, $\phi_{\text{src}} = 0$. The counterclockwise orbit, with $b/M = +6$, rises to a turning point at ($r_{\text{tp}}/M = 2.37$, $\phi_{\text{tp}} = 63^\circ$), then falls back through the event horizon to arrive at the singularity at map angle $\phi_0 = +279^\circ$. The clockwise orbit with $b/M = -6$ crosses the horizon immediately and reaches the singularity at the map angle $\phi_0 = -154^\circ$. The event X locates a falling observer that intercepts the counterclockwise light orbit at ($r_{\text{obs}}/M = 1$, $\phi_{\text{obs}} = 189^\circ$).

- C. An orbit with impact parameter $|b/M| \approx b_{\text{critical}}/M = (27)^{1/2}$ circles at $r \approx 3M$ for a while. Then it “falls off the knife-edge,” either spiraling inward or returning outward to $r/M \gg 1$. In the second case the turning r -coordinate is $r_{\text{tp}}/M \approx 3$, but *where on that circle* is the turning point?

QUERY 7. Infinite impact parameter

- A. From equation (29), find two different conditions that lead to $|b/M| \rightarrow \infty$.
 B. In Figure 3, what horizontal line corresponds to $(M/b)^2 \rightarrow 0$ or $|b/M| \rightarrow \infty$? Point out two places on the graph (one a limiting case) where $(V(r)/M)^2$ reaches this line.

11.6. ■ STARLIGHT ORBIT: FROM STAR TO OBSERVER

375 *Starlight orbit must reach me.*

Which orbit(s) connect(s) the star with the observer?

376 Which light orbit(s) connect(s) a particular star to a given map location near
 377 the black hole? This question is important because sooner or later we want to
 378 predict in what direction one of the many possible inertial observers at that
 379 map location looks to see a particular star. But an observer cannot see light
 380 that does not reach him or her. The central goal of this chapter is to find the
 381 global path of an orbit that connects distant Star X to a given map location
 382 Y, whatever the motion may be of an observer at rest or moving through that
 383 location.

Section 11.6 Starlight Orbit: From Star to Observer 11-17




FIGURE 7 Starlight orbit A with impact parameter $b/M = -8$ moves in a clockwise direction to connect the star at map angle $\phi_\infty = 70.07^\circ$ to observer P located at $(r_{\text{obsP}}/M = 8, \phi_{\text{obsP}} = 0)$. The starlight orbit proceeds to observer Q, crossing outward through the shell at the same $r_{\text{obsQ}}/M = r_{\text{obsP}}/M = 8$ but at a different value ϕ_{obsP} , to be determined.

?

384
385
386

Objection 2. *Ha, gotcha! You say that the observer can be at any coordinate r_{obs} . But inside the event horizon nothing can stand still in global coordinates. Therefore you cannot have an observer at $r_{\text{obs}} < 2M$.*

!

387
388
389
390
391
392

You are correct: No observer can remain at constant r inside the event horizon. However Chapters 6, 7, and 12 describe the rain observer who starts from rest far from the black hole and drops to its center. This rain observer receives starlight even inside the event horizon. To predict the spectacular, ever-changing rain observer's pre-doom panoramas (Chapter 12), we must know which orbit(s) from every star reach(es) her there.

393 The orbit labeled A in Figure 7 connects a distant star to a point with map
394 location $(r_{\text{obs}}/M = 8, \phi_{\text{obs}} = 0)$ where we will later place one of many possible

11-18 Chapter 11 Orbits of Light

395 observers. This figure introduces the **map angle** ϕ_∞ of the distant star. The
 396 subscript infinity, ∞ , reminds us that the star lies far from the black hole.

Map angle ϕ_∞
 to a star

$$\phi_\infty \equiv (\text{map angle to a distant star, this angle measured} \quad (36)$$

counterclockwise from the direction $\phi = 0$)

397 Section 11.7 shows that many orbits—in principle an infinite number of
 398 orbits—from each star arrive at the map location of any observer. How do we
 399 choose which orbit to follow? *Answer:* We discover that there is a single
 400 most-direct orbit between star and observer, an orbit whose spatial path is the
 401 least deflected in map coordinates. We call this the **primary orbit** and give it
 402 most of our attention, often simply calling it “the orbit.”

Primary orbit

403 What primary orbit connects the star at given map angle ϕ_∞ most
 404 directly with the observer at map location $(r_{\text{obs}}, \phi_{\text{obs}} = 0)$? This is an
 405 important question with a complicated answer. So start with an example.

Primary orbit
 between star
 and map location
 of the observer

406 Figure 7 shows the interactive software GRorbits plot of a primary Bounce
 407 Orbit between a star at map angle $\phi_\infty = 70.07^\circ$ and an observer at map
 408 location $(r_{\text{obs}} = 8M, \phi_{\text{obs}} = 0)$. *Result:* The orbit with impact parameter
 409 $b/M = -8$ connects this observer with the star at map angle $\phi_\infty = 70.07^\circ$.

410 The incoming orbit in Figure 7 sweeps clockwise past the observer at
 411 $r/M = 8$, reaches a turning point at smaller r -coordinate, then crosses the
 412 $r/M = 8$ shell a second time, now in an outgoing direction. Two observers
 413 located at *different* points along the same shell can see the same orbit from the
 414 *same* star.

Incoming orbit may
 move out again
 across the same shell.

11.7 ■ INTEGRATE THE STARLIGHT ORBIT

416 *An exact and immediate result*

417 Our goal is to plot $\phi_\infty - \phi_{\text{obs}}$ for starlight as a function of r_{obs} for a given
 418 value of the impact parameter b . To accomplish this, integrate $d\phi/dr$ directly.
 419 Figure 7 shows two cases. Case I: The orbit reaches the observer before the
 420 turning point. Case II: The orbit reaches the observer after the turning point.
 421 Both cases integrate equation (19).

Goal: To plot
 $\phi_\infty - \phi_{\text{obs}}$
 for starlight

$$\phi_\infty - \phi_{\text{obs}} = \int_{r=\infty}^{r_{\text{obs}}} \frac{b}{r^2} F^{-1}(b, r) dr \quad (37)$$

(Case I: observer before turning point)

$$\phi_\infty - \phi_{\text{obs}} = \int_{r=\infty}^{r_{\text{tp}}} \frac{b}{r^2} F^{-1}(b, r) dr + \int_{r_{\text{tp}}}^{r_{\text{obs}}} \frac{b}{r^2} F^{-1}(b, r) dr \quad (38)$$

(Case II: observer after turning point)

422 Figure 8 displays the result of these integrals. The vertical axis “unrolls” the
 423 ϕ -angle.

Section 11.8 Multiple Starlight Orbits from Every Star 11-19



424 **Objection 3.** *How do you carry out these integrals? Function $F(b, r)$ in*
 425 *(16) is complicated; these integrations must be difficult.*



426 **!** Modern numerical methods evaluate these integrals to high accuracy. We
 427 do not pause here to describe these methods.

Plunge Orbit
 has small $|b|$.
 Bounce Orbit
 has large $|b|$.

428 Figure 3 previewed the summary message of Figure 8: An incoming orbit
 429 with small magnitude of $|b|$ plunges through the event horizon to the
 430 singularity. An incoming orbit with a large magnitude of $|b|$ deflects and
 431 returns outward again. An incoming orbit with the particular intermediate
 432 value $\pm b_{\text{critical}}$ circles temporarily at $r = 3M$, then either continues ingoing or
 433 becomes outgoing.



434 **Objection 4.** *You are not telling us the whole story! Orbits in most figures*
 435 *of this chapter have arrows on them. Every arrow tells us the direction of*
 436 *motion of light at that place along the orbit. But motion involves increments*
 437 *in the T -coordinate. Your equations that lead to these figures do not*
 438 *contain global T . Therefore these equations can give us only the curves*
 439 *themselves, without arrows.*



440 **!** Yes and no. Equation (5) defines b as L/E , so the sign of the impact
 441 parameter is the same as the sign of L . This means that the motion of light
 442 is counterclockwise for positive values of b and clockwise for negative
 443 values. So equations (38) and (39) do give us the directions of motion
 444 (arrow directions) simply from the signs of b/M in those equations.
 445 Indeed, these equations do not tell us the map position of each light flash
 446 as a function of the T -coordinate. But we are interested in the plot of a
 447 steady starlight orbit, which does not vary with T .

448 Sample Problems 2 illustrate uses of Figure 8.

449 **Comment 3. Every black hole redirects to every observer multiple orbits**
 450 **from every star.**

451 You can use Figure 8 to find the value b of an orbit that connects *any* distant star
 452 ($-180^\circ < \phi_\infty \leq +180^\circ$) to a map location on *some* circle of *any* r -coordinate
 453 around the black hole. Whoa! Does this mean that the black hole never obscures
 454 any star in the heavens for an observer near it? Yes, and more: The following
 455 section and Figure 10 show that every black hole in the visible Universe redirects
 456 multiple orbits from every single star in the heavens to an observer at every
 457 single map location.

11.8 ■ MULTIPLE STARLIGHT ORBITS FROM EVERY STAR

459 *An infinite number of orbits that appear fainter and fainter to an observer.*

One star:
 Infinite images?

460 It is remarkable that every map location near a black hole receives multiple
 461 orbits—in principle an infinite number of orbits—from a single star, and thus

11-20 Chapter 11 Orbits of Light




FIGURE 8 Difference in map angles between a distant star and the observer at map location $(r_{\text{obs}}/M, \phi_{\text{obs}})$ derived for an orbit of impact parameter b/M from that star. To reduce clutter, we define $b^* \equiv b/M$. Arrows on the curves tell whether the starlight is incoming or outgoing; at a turning point the orbit changes from incoming to outgoing.

462 from every star in the heavens. Figure 9 replots the primary orbit of Figure 7
 463 and adds two additional orbits, called **higher-order orbits** from the same
 464 star. By trial and error, the interactive software program GRorbits finds values
 465 $b/M = +5.4600$ and $b/M = -5.2180$ for these additional orbits from the same
 466 star.

Higher-order
orbits

467 In Figure 9, the higher-order orbit with $b/M = +5.4600$ moves around the
 468 black hole counterclockwise and approaches the map location
 469 $(r/M = 8, \phi = 0)$ from below. This orbit lacks 70.07° of making a complete
 470 circuit around the black hole. Therefore the *total* angle to the same star is
 471 $\phi_\infty = -(360^\circ - 70.07^\circ) = -289.93^\circ$.

472 The next higher-order orbit with $b/M = -5.2180$ moves around the black
 473 hole clockwise and approaches the map location $(r/M = 8, \phi = 0)$ from above.
 474 This orbit makes a complete circuit around the black hole, *plus* 70.07° , for a
 475 total of 430.07° . Therefore the *total* angle to the same star is
 476 $\phi_\infty = +(360^\circ + 70.07^\circ) = +430.07^\circ$.

Each observer
receives many
orbits from
every star.

477 Figure 10 extends the vertical scale of Figure 8 to show orbits with
 478 b -values close to the critical value that circle several times around the black
 479 hole before they either escape outward or plunge on inward. The upward and
 480 downward vertical scales in Figure 10 extend indefinitely, leading to more and
 481 more orbits with b -values on either side of $b_{\text{critical}}/M = (27)^{1/2} = 5.196152\dots$
 482 *Conclusion:* An observer at each r -coordinate r_{obs} receives multiple orbits—in
 483 principle an infinite number of orbits—from every star in the heavens.

Section 11.8 Multiple Starlight Orbits from Every Star 11-21

Sample Problems 1. Orbits that reach $r/M = 3$

Think of orbits with different b -values that reach the observer map location at $(r_{\text{obs}}/M = 3, \phi_{\text{obs}} = 0)$. Use Figure 8 to provide approximate answer the following questions.

- A. What is the b -value of the orbit that comes from the star at map angle $\phi_{\infty} = +60^\circ$? **Solution A:** Look at the vertical dashed line at $r_{\text{obs}}/M = 3$. This line intersects with the horizontal line $\phi_{\infty} = +60^\circ$ very close to the curve $b/M = -3$, at the point marked G. So this is the b -value of the Plunge Orbit that connects the star at map angle $\phi_{\infty} = +60^\circ$ with the observer at $(r_{\text{obs}}/M = 3, \phi_{\text{obs}} = 0)$.
- B. What is the b -value of the orbit that comes from the star at map angle $\phi_{\infty} = +90^\circ$? **Solution B:** The vertical dashed line at $r_{\text{obs}}/M = 3$ intersects the horizontal line $\phi_{\infty} = +90^\circ$ very close to the Plunge Orbit $b/M = -4$.
- C. What is the b -value of the orbit that comes from the star at map angle $\phi_{\infty} = +30^\circ$? **Solution C:** The vertical dashed line $r_{\text{obs}}/M = 3$ intersects with the horizontal line $\phi_{\infty} = +30^\circ$ about six-tenths of the separation between

the curves $b/M = -1$ and $b/M = -2$. Therefore the Plunge Orbit with $b \approx -1.6$ connects the star at map angle $\phi_{\infty} = +30^\circ$ with the map location $(r_{\text{obs}}/M = 3, \phi_{\text{obs}} = 0)$.

- D. What is the b -value of the orbit that comes from the star at negative map angle $\phi_{\infty} = -90^\circ$? **Solution D:** The vertical dashed line $r_{\text{obs}}/M = 3$ intersects the horizontal line $\phi_{\infty} = -90^\circ$ very close to the curve $b/M = +4$. The positive b -value means that the orbit moves counterclockwise around the black hole.
- E. an orbit comes from the opposite side of the black hole, at $\phi_{\infty} = 180^\circ$. What is the b -value of this orbit? **Solution E:** Both $\phi_{\infty} = +180^\circ$ and $\phi_{\infty} = -180^\circ$ are map angles to a star on the other side of the black hole. The vertical dashed line $r_{\text{obs}}/M = 3$ intersects the horizontal lines $\phi_{\infty} = \pm 180^\circ$ approximately half way between $b/M = \pm 5$ and $b/M = \pm(27)^{1/2} = \pm 5.196$. Therefore the b -values of these two Plunge Orbits are approximately $b \approx \pm 5.1$. *Optional:* Sketch this orbit.

Sample Problems 2. Orbits from a single star that reach observers at different r -coordinates

Orbits with different b -values from the star at map angle $\phi_{\infty} = +60^\circ$ reach observers at different r -coordinates along the line $\phi = 0$. What are these b -values at r -coordinates $r_{\text{obs}}/M = 12, 8, 4, 2, \text{ and } 1$? In each case say whether the orbit is a Plunge Orbit, a Bounce Orbit, or a Trapped Orbit.

Solution: All of the orbits are from a star; therefore none of them can be a Trapped Orbit. In Figure 8, look at the intersections of horizontal line $\phi_{\infty} = +60^\circ$ with vertical lines

at these different r -coordinates. We estimate the b -values to one decimal place.

- At $r_{\text{obs}}/M = 12, b/M \approx -10.9$, the point marked F in the figure; a Bounce Orbit
- At $r_{\text{obs}}/M = 8, b/M \approx -7.3$, a Bounce Orbit
- At $r_{\text{obs}}/M = 4, b/M \approx -3.8$, a Plunge Orbit
- At $r_{\text{obs}}/M = 2, b/M \approx -2.0$, a Plunge Orbit
- At $r_{\text{obs}}/M = 1, b/M \approx -1.2$, a Plunge Orbit

484 Look at the little square white boxes on the vertical line at $r/M = 8$ in
 485 Figure 10. Three of the little white boxes on the vertical line at $r/M = 8$
 486 correspond to the three starlight orbits displayed in Figure 9. Other little boxes
 487 represent more of the multiple higher-order orbits between this star and this
 488 observer. Each little box is offset vertically by $\pm 360^\circ$ from its nearest neighbor.

489

QUERY 8. *Optional:* Classify primary and higher-order orbits from a star.

11-22 Chapter 11 Orbits of Light




FIGURE 9 Three of the infinite number of orbits of light that, in principle, arrive at the same observer from a single star. For the primary orbit with $b/M = -8$, the star angle is $\phi_\infty = 70.07^\circ$ (as in Figure 7). For the second orbit, with $b/M = +5.4600$, the star angle (dashed arc) is $\phi_\infty = -(360^\circ - 70.07^\circ) = -289.93^\circ$. For the third orbit, with $b/M = -5.2180$, the star angle (angle-arc not shown) is $\phi_\infty = (360^\circ + 70.07^\circ) = +430.07^\circ$. All three orbits come from the same star, but the observer sees three different images in three different directions.

Classify the primary and higher-order starlight orbit as a Plunge Orbit or a Bounce Orbit. Figure 10 may be useful. *Reminder:* This analysis says nothing about the state of motion of the observer at that map location: he may be at rest there; she may dive or orbit past that map location.

- A. Show that for every observer inside $r/M = 3$, all starlight orbits are Plunge Orbits.
- B. Show that for every observer outside $r/M = 3$, starlight orbits are either Plunge Orbits or Bounce Orbits.
- C. At any $r/M > 3$, what is the value of b/M that divides Plunge Orbits from Bounce Orbits?
- D. Find an equation for the maximum magnitude of the impact parameter b/M of a Bounce Orbit that an observer on the shell of a given r -coordinate $r/M > 3$ can see?
- E. Show that for every observer at $r/M > 3$, every higher-order orbit is an outgoing Bounce Orbit.
- F. Can a primary or higher-order starlight orbit be a Trapped Orbit? Explain your answer.

Section 11.8 Multiple Starlight Orbits from Every Star 11-23




FIGURE 10 Expanded vertical scale for starlight orbits of Figure 8. The observer is at map location $(r_{obs}/M, \phi_{obs})$. *New feature of this plot:* Orbits with $b^* \approx \pm b_{critical}/M$ follow the vertical line at $r/M = 3$ (they circulate at $r/M = 3$) before they either return to $r/M \gg 1$ or plunge into the black hole. *Result:* Multiple orbits—in principle an infinite number of orbits—from every star arrive at each observer, cross every possible vertical line in the figure. *Example:* Three of the little white boxes on the vertical line at $r/M = 8$ correspond to the three starlight orbits displayed in Figure 9.

Higher-order orbits
have fainter,
smeared images.

503 Higher-order orbits that go around the black hole more and more times
504 are less and less intense when they arrive at the observer. There is always
505 *some* spread in the orbit, so the more times an orbit circles the black hole, the
506 more it spreads out transverse to its direction of motion and the smaller the

11-24 Chapter 11 Orbits of Light




FIGURE 11 Forbidden region for light. Near the non-spinning black hole, this forbidden region separates our world, above the forbidden region, from another world, below the forbidden region.

507 fraction of photons in the initial orbit that enter the detector at the final map
 508 location. Chapter 12 shows that the shell observer also sees higher-order orbits
 509 bunched closer and closer together in the observed direction. *Overall result:*
 510 Higher and higher order orbits lead to images that get fainter and fainter and
 511 smear into one another. As a result, an observer sees separately only a few of
 512 the infinite number of orbits that, in principle, arrive from each star.

513 Strange results follow from equation (24), which expresses $(dr/dT)^2$ in
 514 terms of the difference $(M/b)^2 - (V(r)/M)^2$. Differentials dr and dT are both
 515 real, so dr/dT must be real. In other words $(dr/dT)^2$ must be positive.

516 *Conclusion:* $(M/b)^2 - (V(r)/M)^2$ must be positive. A consequence of this
 517 condition is that either $M/b > +V(r)/M$ or $M/b < -V(r)/M$. The result is a
 518 forbidden region where light cannot exist, as shown in Figure 11. Compare
 519 corresponding Figure 5 in Section 8.4 for the stone and review the text that
 520 accompanies that figure. Near the black hole the forbidden region for light
 521 separates our world (above the forbidden region) from another world (below
 522 the forbidden region). We can move between these worlds only by entering and
 523 then exiting the event horizon—not possible for a non-spinning black hole.

524 However, we will find that for the spinning black hole a trip from the
 525 corresponding upper region to the corresponding lower region may be possible.
 526 John Archibald Wheeler’s radical conservatism says, “Follow the equations
 527 wherever they lead, no matter how strange the result.”

Two worlds,
 separated for the
 non-spinning
 black hole

11.9 ■ EXERCISES

529 **Note:** In the exercises the word *approximately* means that the requested
 530 number may be estimated from a figure in this chapter.

531 **1. Thought question: Shadow of a Black Hole?**

532 According to legend, a vampire has no reflection in a mirror and casts no
 533 shadow. When illuminated from one side by a distant incoming flat wave, does
 534 a black hole cast a shadow on the other side? Think of a possible shadow on a
 535 flat plane located far away from the black hole where spacetime is flat.

536 **2. Values of b for orbits that arrive at $r_{\text{obs}}/M = 6$.**

537 Repeat parts A through E of Sample Problems 2 for orbits that reach the
 538 observer at map location ($r_{\text{obs}}/M = 6, \phi_{\text{obs}} = 0$). Classify each orbit as
 539 incoming, outgoing, or tangential.

540 **3. Orbits that reach observers at different r -coordinates from the star at map
 541 angle $\phi_{\infty} = -120^\circ$.**

542 Repeat Sample Problems 2 for a star at map angle $\phi_{\infty} = -120^\circ$.

543 **4. The visual size of a black hole**

544 Figure 10 shows the b -values of beams that escape or are captured by the
 545 black hole. The smallest b -value of a beam that can escape is
 546 $|b_{\text{critical}}| = (27)^{1/2}M$. Some light from every star circles temporarily on this
 547 unstable orbit at $r = 3M$. Because this is a knife-edge orbit, it continually
 548 sheds light beams that “fall off” to move either inward or outward.




FIGURE 12 Schematic diagram showing the visual size of the black hole Sagittarius A* located at the center of our galaxy, assumed (incorrectly) to be non-spinning. The text shows that all possible parallel straight beams form a three-dimensional cylinder directed toward the observer on Earth.

11-26 Chapter 11 Orbits of Light




FIGURE 13 Critical beams from Sgr A* form a long cone as seen from Earth

549 Consider outward light beams that enter the eye of a distant observer on
 550 Earth. Figure 12 shows two such beams on one $[r, \phi]$ slice through the center of
 551 the black hole. But the same distant observer sees a similar pair of beams that
 552 lie on each of an infinite number of similar slices rotated around the r -axis in
 553 Figure 12. The resulting set of beams form a cylinder observed by the Earth
 554 observer.

555 To speak more carefully, the beams we see on Earth do not move *exactly*
 556 on a cylinder, but rather on a very long cone with its apex at the Earth
 557 (Figure 13). As a result, we on Earth see the black hole as a ring. What angle
 558 does this ring subtend at our eye on Earth?

559 Answer this question for the monster black hole called Sagittarius A*
 560 (abbreviation: SgrA*) with mass $M_{\text{SgrA}} \approx 4 \times 10^6 M_{\text{Sun}}$ that lies at the center
 561 of our galaxy, about 26 000 light-year from Earth. Label this distance r_{Earth} .
 562 Assume (incorrectly) that SgrA* is a nonspinning black hole. Derive and
 563 justify an expression for the angular size θ_{Earth} of this black hole observed
 564 from Earth. (An exercise in Chapter 20 carries out a more realistic analysis
 565 that takes account of the spin of this black hole.)

- 566 A. From Figure 13, derive the following expression for the very small angle
 567 θ_{Earth} .

$$\theta_{\text{Earth}} \approx \frac{2(27)^{1/2} M_{\text{SgrA}}}{r_{\text{Earth}}} \quad (r \gg M_{\text{SgrA}}) \quad (39)$$

- 568 B. Insert into (39) values for M_{SgrA} and Earth's r -coordinate separation
 569 from the black hole of r_{Earth} light years. The following are results to
 570 one significant digit. Find each result to two significant digits:

$$\begin{aligned} \theta_{\text{Earth}} &\approx 2 \times 10^{-10} && \text{radian} && (40) \\ &\approx 1 \times 10^{-8} && \text{degree} \\ &\approx 5 \times 10^{-5} && \text{arcsecond} \\ &\approx 50 && \text{microarcseconds} \end{aligned}$$

571 **Comment 4. Microwaves, not visible light**

572 Dust between Earth and the spinning black hole at the center of our galaxy
 573 absorbs visible light. Microwaves pass through this dust, so our detectors on
 574 Earth are microwave dishes distributed over the surface of Earth.

575 **5. The “incoming map floodlight”**

576 Define an **incoming map floodlight** as a lamp at a given r -coordinate
 577 r_{inlamp} that emits all light beams that are ingoing at that r —that is, all beams
 578 with a negative r -coordinate differential, $dr < 0$.

- 579 A. An incoming map floodlight at $r_{\text{inlamp}}/M = 12$ emits light that might
 580 have come from stars with approximately what range of map angles
 581 ϕ_∞ ?
- 582 B. An incoming map floodlight at $r_{\text{inlamp}}/M = 6$ emits light that might
 583 have come from stars with approximately what range of map angles
 584 ϕ_∞ ?
- 585 C. An incoming map floodlight at $r_{\text{inlamp}}/M = 3$ emits light that may have
 586 come from stars with approximately what range of map angles ϕ_∞ ?
- 587 D. An incoming map floodlight at $r_{\text{inlamp}}/M = 1$ emits light that may have
 588 come from stars with approximately what range of map angles ϕ_∞ ?
- 589 E. Can the incoming map floodlight at $r_{\text{inlamp}}/M = 6$ be at rest in global
 590 coordinates? Can the incoming map floodlight at $r_{\text{inlamp}}/M = 1$ be at
 591 rest in global coordinates?

592 **6. The “outgoing map floodlight”**

593 Define an **outgoing map floodlight** as a lamp at a given r -coordinate,
 594 r_{outlamp} , that emits all light beams that are outgoing at that
 595 r -coordinate—that is, all beams with a positive r -coordinate differential,
 596 $dr > 0$.

- 597 A. An outgoing map floodlight at $r_{\text{outlamp}}/M = 8$ emits light that might
 598 have come from stars with approximately what range of map angles
 599 ϕ_∞ ?
- 600 B. An outgoing map floodlight at $r_{\text{outlamp}}/M = 5$ emits light that may
 601 have come from stars with approximately what range of map angles
 602 ϕ_∞ ?
- 603 C. An outgoing map floodlight at $r_{\text{outlamp}}/M = 3$ emits light that may
 604 have come from stars with approximately what range of map angles
 605 ϕ_∞ ?
- 606 D. Is there a range of r -coordinates in which the outgoing map floodlight
 607 is useless? *Hint:* look at Figure 10.

608 **7. Newton’s plot of map angle difference.**

609 Make a *rough* sketch (don’t sweat the details) of Figure 8 for orbits of light in
 610 Newtonian mechanics, in which spacetime is flat around the center of
 611 attraction and light is fast particle. What “Newtonian assumptions” do you

11-28 Chapter 11 Orbits of Light

612 make about the path of light under this attraction? (We have no record that
613 Newton himself made any prediction about the effect of his “gravitational
614 force” on the orbits of light.)

11.10 ■ REFERENCES

616 Initial quotes:

617 Egyptian creation quote from

618 [M=http://www.aldokkan.com/religion/creation.htm/](http://www.aldokkan.com/religion/creation.htm/)

619 Tuamotuan creation quote from *The Myths of Creation* by Charles H. Long,
620 George Braziller, New York, 1963, pages 173 and 179.

621 Inuit creation quote from *The Power of Stars—How Celestial Observations*
622 *Have Shaped Civilization* by Bryan E. Penprase, New York, Springer 2011,
623 page 97.

624 The interactive GRorbits program that plots orbits of light is available at
625 website <http://stuleja.org/grorbits/>

626 Download File Name: Ch11OrbitsOfLight170511v1.pdf